



Application of the subordination principle to the multivalent harmonic mappings with shear construction method

Yaşar Polatoğlu, Emel Yavuz Duman*, H. Esra Özkan

Department of Mathematics and Computer Science, İstanbul Kültür University, 34156 İstanbul, Turkey

ARTICLE INFO

Article history:

Received 7 September 2010

Accepted 23 December 2010

Keywords:

Harmonic functions
Convex in one direction
Growth theorem
Distortion theorem

ABSTRACT

The harmonic function in the open unit disc $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ can be written as a sum of an analytic and an anti-analytic function, $f = h(z) + g(z)$, where $h(z)$ and $g(z)$ are analytic functions in $\mathbb{D}$, and are called the analytic part and co-analytic part of f , respectively.

One of the most important questions in the study of the classes of such functions is related to bounds on the modulus of functions (growth) or modulus of the derivative (distortion), because the growth theorem and distortion theorem give the compactness of the classes of these functions. In this paper we consider both of these questions with the shear construction method.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Let $\mathcal{U}$ be a simply connected domain in the complex plane $\mathbb{C}$. A harmonic function f has the representation $f = h(z) + \overline{g(z)}$, where $h(z)$ and $g(z)$ are analytic in $\mathcal{U}$ and are called the analytic and co-analytic part of f , respectively. Let $h(z) = z^m + a_{m+1}z^{m+1} + a_{m+2}z^{m+2} + \dots$ and $g(z) = b_m z^m + b_{m+1}z^{m+1} + b_{m+2}z^{m+2} + \dots$ ($m \in \mathbb{N}$) be analytic functions in the open unit disc $\mathbb{D}$. The Jacobian of the mapping f , denoted by $J_f(z)$, and can be computed by $J_f(z) = |h'(z)|^2 - |g'(z)|^2$. If $J_f(z) = |h'(z)|^2 - |g'(z)|^2 > 0$, then f is called a sense-preserving multivalent harmonic function. The class of all sense-preserving multivalent harmonic functions with $|b_m| < 1$ is denoted by $\mathcal{H}_m(m)$, and the class of all sense-preserving multivalent harmonic functions with $b_m = 0$ is denoted by $\mathcal{H}_m^0(m)$. For convenience, we will investigate sense-preserving harmonic functions, that is, functions for which $J_f(z) > 0$. If $J_f(z) < 0$, then $\bar{f}$ is sense-preserving. The second analytic dilatation of a harmonic function is given by $w(z) = g'(z)/h'(z)$. We also note that if f is locally univalent and sense-preserving, then $|w(z)| < 1$ for every $z \in \mathbb{D}$.

In this paper we examine the class of functions that are convex in the direction of real axis. The shear construction is essential to the present work as it allows one to study harmonic functions through their related analytic functions ([1] Hengartner and Shober). The shear construction produces a univalent harmonic function that maps $\mathbb{D}$ to a region that is convex in the direction of the real axis. This construction relies on the following theorem of Clunie and Sheil-Small.

Theorem 1.1 ([2]). *A harmonic function $f = h(z) + \overline{g(z)}$ locally univalent in $\mathbb{D}$ is a univalent mapping of $\mathbb{D}$ onto a domain convex in the direction of the real axis if and only if $h(z) - g(z)$ is a conformal univalent mapping of $\mathbb{D}$ onto a domain convex in the direction of the real axis.*

Hengartner and Shober [1] studied analytic functions $\psi(z)$ that are convex in the direction of the imaginary axis. They used a normalization which requires, in essence, that the right and left extremes of $\psi(\mathbb{D})$ be the images of 1 and -1 . This

* Corresponding author.

E-mail addresses: y.polatoglu@iku.edu.tr (Y. Polatoğlu), e.yavuz@iku.edu.tr (E.Y. Duman), e.ozkan@iku.edu.tr (H. Esra Özkan).

normalization is as follows: there exist points z'_n converging to $z = 1$ and z''_n converging to $z = -1$ such that

$$\begin{aligned}\lim_{n \rightarrow \infty} \operatorname{Re}\{\psi(z'_n)\} &= \sup_{|z| < 1} \operatorname{Re}\{\psi(z)\}, \\ \lim_{n \rightarrow \infty} \operatorname{Re}\{\psi(z''_n)\} &= \inf_{|z| < 1} \operatorname{Re}\{\psi(z)\}.\end{aligned}\quad (1.1)$$

If $\mathcal{CLA}$ is the class of functions on the domains, D , that are convex in the direction of the imaginary axis and admit a mapping $\psi(z)$ so that $\psi(\mathbb{D}) = D$ and $\psi(z)$ satisfies the normalization (1.1), then we have the following result:

Theorem 1.2 ([1]). Suppose that $\psi(z)$ is analytic and non-constant for $|z| < 1$. Then we have $\operatorname{Re}[(1 - z^2)\psi'(z)] \geq 0$ for $|z| < 1$ if and only if

- (i) $\psi(z)$ is univalent on $\mathbb{D}$,
- (ii) $\psi(\mathbb{D}) \in \mathcal{CLA}$, and
- (iii) $\psi(z)$ is normalized by (1.1).

Remark 1.3 ([1]). The condition $\operatorname{Re}[(1 - z^2)\psi'(z)] \geq 0$ (for $|z| < 1$), has an elementary geometric interpretation. If we parametrize the line segment and circular arcs γ_t , $-\pi/2 < t < \pi/2$, joining $z = -1$ to $z = 1$ in the unit disc by

$$\gamma_t : z = z(s) = \frac{e^{s+it} - 1}{e^{s+it} + 1}, \quad -\infty < s < \infty,$$

then one easily verifies that

$$\frac{d}{ds} \operatorname{Re}[\psi(z(s))] = 2\operatorname{Re}[(1 - z^2(s))\psi'(z(s))].$$

Consequently, the condition $\operatorname{Re}[(1 - z^2)\psi'(z)] \geq 0$ (for $|z| < 1$) is equivalent to the property that the circular arcs γ_t are mapped onto analytic arcs which may be represented as functions $v = v(u)$. It follows that $\psi(z)$ has the normalization (1.1). Furthermore, since the region bounded by $\psi(\gamma_t) \cup \psi(1) \cup \psi(-1)$ is convex in the v -direction for every $-\pi/2 < t < \pi/2$, we find that $\psi(|z| < 1)$ is also convex in the v -direction.

Remark 1.4 ([1]). An analytic function $\psi(z)$ is close-to-convex if there exists a convex mapping $s(z)$ such that $\operatorname{Re}[\psi'(z)/s'(z)] > 0$ for $|z| < 1$. Functions satisfying $\operatorname{Re}[(1 - z^2)\psi'(z)] > 0$ are special close-to-convex functions associated with $s(z) = \frac{1}{2} \log[(1 + z)/(1 - z)]$. Kaplan [3] has shown that close-to-convex functions, hence functions satisfying $\operatorname{Re}[(1 - z^2)\psi'(z)] > 0$, are univalent. The geometric interpretation of Remark 1.3 could also be used to show that functions satisfying $\operatorname{Re}[(1 - z^2)\psi'(z)] > 0$ are univalent.

Remark 1.5 ([4]). We also note that Theorem 1.1 has a natural generalization when f is convex in the direction α . In that situation $e^{-i\alpha}f$ and $\varphi(z) = e^{-i\alpha}h(z) - e^{i\alpha}g(z)$ are convex in the direction of the real axis, hence the function $h(z) - e^{i2\alpha}g(z)$ is convex in the direction α . In particular, we can use this construction when $\alpha = \pi/2$ to construct functions that are convex in the direction of the imaginary axis. At the same time, to be able to use this result for functions that are convex in the direction of the real axis, let us consider the following situation: suppose that $\varphi(z)$ is a function that is analytic and convex in the direction of the real axis. Furthermore, suppose that φ is normalized by the following. Let there exist points z'_n converging to $z = e^{i\alpha}$ and z''_n converging to $z = e^{i(\alpha+\pi)}$ such that

$$\begin{aligned}\lim_{n \rightarrow \infty} \operatorname{Im}\{\varphi(z'_n)\} &= \sup_{|z| < 1} \operatorname{Im}\{\varphi(z)\}, \\ \lim_{n \rightarrow \infty} \operatorname{Im}\{\varphi(z''_n)\} &= \inf_{|z| < 1} \operatorname{Im}\{\varphi(z)\}.\end{aligned}\quad (1.2)$$

Consequently, if $\psi(z)$ satisfies (1.1), then $\varphi(z) = i\psi(e^{-i\alpha}z)$ satisfies (1.2).

Finally, let Ω be the family of functions $\phi(z)$ which are regular in $\mathbb{D}$ and satisfying the conditions $\phi(0) = 0$, $|\phi(z)| < 1$ for every $z \in \mathbb{D}$. Denote by $\mathcal{P}(m)$ (with m being a positive integer) the family of functions $p(z) = m + p_1z + p_2z^2 + \dots$ which are regular in $\mathbb{D}$ and satisfying the conditions $p(0) = m$, $\operatorname{Re}p(z) > 0$ for all $z \in \mathbb{D}$, and such that $p(z)$ is in $\mathcal{P}(m)$ if and only if

$$p(z) = m \frac{1 + \phi(z)}{1 - \phi(z)} \quad (1.3)$$

for some function $\phi(z) \in \Omega$ and every $z \in \mathbb{D}$. Let $F(z) = z + \alpha_2z^2 + \dots$ and $G(z) = z + \beta_2z^2 + \dots$ be analytic functions in $\mathbb{D}$, if there exists a function $\phi(z) \in \Omega$ such that $F(z) = G(\phi(z))$ for all $z \in \mathbb{D}$, then we say that $F(z)$ subordinate to $G(z)$ and we write $F(z) \prec G(z)$. We also note that if $F(z) \prec G(z)$, then $F(\mathbb{D}) \subset G(\mathbb{D})$.

Denote by $\mathcal{S}_{\mathcal{H}}^0(\mathcal{C}(m))$ the class of all m -valent harmonic functions convex in the direction of real axis. In this paper we will give growth and distortion theorems for the class $\mathcal{S}_{\mathcal{H}}^0(\mathcal{C}(m))$.

2. Main results

Lemma 2.1. Let $\varphi(z) = z^m + c_{m+1}z^{m+1} + c_{m+2}z^{m+2} + \dots$ be analytic in $\mathbb{D}$. If $\varphi(z)$ satisfies the condition $\operatorname{Re} \left[(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} \right] > 0$, then $\varphi(z)$ is an m -valent close-to-convex function and

$$\frac{mr^{m-1}(1-r)}{(1+r^2)^m(1+r)} \leq |\varphi'(z)| \leq \frac{mr^{m-1}}{(1+r)^{m-1}(1-r)^{m+1}} \quad (2.1)$$

for all $|z| = r < 1$.

Proof. Let us consider the function $s(z) = \int_0^z \frac{\zeta^{m-1}}{(1-\zeta^2)^m} d\zeta$. Since

$$1 + z \frac{s''(z)}{s'(z)} = m \frac{1+z^2}{1-z^2} \Rightarrow \operatorname{Re} \left(1 + z \frac{s''(z)}{s'(z)} \right) > 0$$

and

$$\operatorname{Re} \left(\frac{\varphi'(z)}{s'(z)} \right) = \operatorname{Re} \left[(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} \right] > 0,$$

then $\varphi(z)$ is a close-to-convex function for all $z \in \mathbb{D}$. On the other hand, since

$$\operatorname{Re} \left[(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} \right] > 0, \quad \left[(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} \right] \Big|_{z=0} = m,$$

$$p(z) \in \mathcal{P}(m) \Leftrightarrow p(z) = m \frac{1 + \phi(z)}{1 - \phi(z)} \Leftrightarrow \phi(z) = \frac{p(z) - m}{p(z) + m},$$

for some $\phi(z) \in \Omega$, the function

$$\phi(z) = \frac{(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} - m}{(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} + m}$$

satisfies the conditions of the Schwarz lemma, whence $|\phi(z)| \leq r$. Therefore we have

$$\left| \frac{(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} - m}{(1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} + m} \right| \leq r \Rightarrow \left| (1 - z^2)^m \frac{\varphi'(z)}{z^{m-1}} - m \frac{1 + r^2}{1 - r^2} \right| \leq \frac{2mr}{1 - r^2}. \quad (2.2)$$

After straightforward calculations we obtain (2.1). $\square$

Theorem 2.2. Let $f = h(z) + \overline{g(z)}$ be an element of $\mathcal{S}_{\mathcal{H}}^0(m)$. Then

$$\frac{mr^{m-1}(1-r)}{(1+r^2)^m(1+r)^2} \leq |f_z| \leq \frac{mr^{m-1}}{(1+r)^{m-1}(1-r)^{m+2}} \quad (2.3)$$

and

$$\frac{|w(z)|mr^{m-1}(1-r)}{(1+r^2)^m(1+r)^2} \leq |f_{\bar{z}}| \leq \frac{mr^m}{(1+r)^{m-1}(1-r)^{m+2}} \quad (2.4)$$

for all $|z| = r < 1$, where $w = g'/h'$ is the second analytic dilatation of f .

Proof. Using Theorems 1.1 and 1.2, Remarks 1.3–1.5, we take $\varphi(z) = h(z) - g(z) \Rightarrow \varphi'(z) = h'(z) - g'(z)$. On the other hand, we have $g'(z) = w(z)h'(z)$, from which it follows that

$$h'(z) = f_z = \frac{\varphi'(z)}{1 - w(z)} \Rightarrow h(z) = \int_0^z \frac{\varphi'(\zeta)}{1 - w(\zeta)} d\zeta, \quad (2.5)$$

$$g'(z) = \bar{f}_{\bar{z}} = \frac{\varphi'(z)w(z)}{1 - w(z)} \Rightarrow g(z) = \int_0^z \frac{\varphi'(\zeta)w(\zeta)}{1 - w(\zeta)} d\zeta, \quad (2.6)$$

and

$$\begin{aligned} g(z) &= \int_0^z \frac{\varphi'(\zeta)w(\zeta)}{1 - w(\zeta)} d\zeta = \int_0^z \frac{\varphi'(\zeta)w(\zeta) - \varphi'(\zeta) + \varphi'(\zeta)}{1 - w(\zeta)} d\zeta \\ &= \int_0^z \frac{\varphi'(\zeta)}{1 - w(\zeta)} d\zeta - \int_0^z \varphi'(\zeta) d\zeta \end{aligned}$$

so that

$$\begin{aligned} f &= h(z) + \overline{g(z)} = \int_0^z \frac{\varphi'(\zeta)}{1-w(\zeta)} d\zeta + \overline{\int_0^z \frac{\varphi'(\zeta)}{1-w(\zeta)} d\zeta} - \int_0^z \varphi'(\zeta) d\zeta \\ &= \operatorname{Re} \left(\int_0^z \frac{2\varphi'(\zeta)}{1-w(\zeta)} d\zeta \right) - \overline{\varphi(z)}, \end{aligned} \quad (2.7)$$

$$\frac{|\varphi'(z)|}{1+r} \leq |f_z| \leq \frac{|\varphi'(z)|}{1-r}, \quad (2.8)$$

and

$$\frac{|w(z)||\varphi'(z)|}{1+r} \leq |f_{\bar{z}}| \leq \frac{r|\varphi'(z)|}{1-r}. \quad (2.9)$$

Using Lemma 2.1 in (2.8) and (2.9), we get (2.3) and (2.4). We note that the inequalities are sharp because the extremal functions can be found as

$$(1-z^2)^m \frac{\varphi'(z)}{z^{m-1}} = m \frac{1+z}{1-z},$$

(i.e., $p(z) \in \mathcal{P}(m)$ then $\frac{1}{p(z)} \in \mathcal{P}(m)$), which apply to (2.7). $\square$

Corollary 2.3. If we take $m = 1$ we obtain

$$\begin{aligned} \frac{1-r}{(1+r^2)(1+r)^2} &\leq |f_z| \leq \frac{1}{(1-r)^3}, \\ \frac{|w(z)|(1-r)}{(1+r^2)(1+r)^2} &\leq |f_{\bar{z}}| \leq \frac{r}{(1-r)^3}. \end{aligned}$$

These inequalities were obtained by Schambroek [4].

Corollary 2.4. Let $f = h(z) + \overline{g(z)}$ be in $\mathcal{H}_\mathcal{H}^0(m)$. Then

$$\frac{m^2 r^{2(m-1)} (1-r)^3}{(1+r^2)^{2m} (1+r)^3} \leq J_{f(z)} \leq \frac{m^2 r^{2(m-1)} (1+r^2)}{(1+r)^{2(m-1)} (1-r)^{2(m+2)}} \quad (2.10)$$

for every $|z| = r < 1$.

Proof. Since $w(z)$ satisfies the conditions of Schwarz lemma, then we have $|w(z)| \leq r$ i.e.,

$$1-r^2 \leq 1-|w(z)|^2 \leq 1+r^2. \quad (2.11)$$

On the other hand, we have

$$J_{f(z)} = |f_z|^2 - |f_{\bar{z}}|^2 = |h'(z)|^2 - |g'(z)|^2 = |h'(z)|^2 (1-|w(z)|^2). \quad (2.12)$$

Using (2.11), (2.12) and Theorem 2.2, we get (2.10) after simple calculations. $\square$

Corollary 2.5. If we take $m = 1$ in Corollary 2.4 we have

$$\frac{(1-r)^3}{(1+r^2)(1+r)^3} \leq J_{f(z)} \leq \frac{1+r^2}{(1-r)^6}.$$

Corollary 2.6. If $f = h(z) + \overline{g(z)} \in \mathcal{H}_\mathcal{H}^0(m)$, then

$$|f| \leq \frac{r^m}{1+m} \left[4mr {}_2F_1(1+m, 2+m, m, 2+m, r, -r) + (1+m) {}_2F_1\left(\frac{m}{2}, m, 1+\frac{m}{2}, r^2\right) \right] \quad (|z| = r < 1), \quad (2.13)$$

where ${}_2F_1(1+m, 2+m, m, 2+m, r, -r)$ is the Appell hypergeometric function, and ${}_2F_1\left(\frac{m}{2}, m, 1+\frac{m}{2}, r^2\right)$ is the Gauss hypergeometric function.

Proof. Since

$$(|h'(z)| - |g'(z)|)|dz| \leq |f| \leq (|h'(z)| + |g'(z)|)|dz|,$$

using Theorem 2.2 we can write

$$|f| \leq \int_0^r \frac{m\rho^{m-1}}{(1+\rho)^{m-2}(1-\rho)^{m+2}} d\rho,$$

which gives (2.13). $\square$

Corollary 2.7. *If we take $m = 1$ we have*

$$|f| \leq \int_0^r \frac{d\rho}{(1-\rho)^3} + \int_0^r \frac{\rho d\rho}{(1-\rho)^3} = \frac{r}{(1-r)^2}.$$

This growth was obtained by Schaubroeck [4].

References

- [1] W. Hengartner, G. Schober, On schlicht mappings to domains convex in one direction, *Comment. Math. Helv.* 45 (1970) 303–314.
- [2] J. Clunie, T. Sheil-Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. AI* 9 (1984) 3–25.
- [3] W. Kaplan, Close-to-convex schlicht functions, *Michigan Math. J.* 1 (1952) 169–185.
- [4] L.E. Schaubroeck, Growth, distortion and coefficient bounds for plane harmonic mappings convex in one direction, *Rocky Mountain J. Math.* 31 (2) (2001) 625–639.